

Universal relation between longitudinal and transverse conductivities in quantum Hall effect

Igor Ruzin¹ and Shechao Feng^{1,2}

1. Department of Physics, University of California, Los Angeles, CA 90024-1547, USA;

2. Department of Physics, Hong Kong University of Science and Technology,
Clear Water Bay Road, Kowloon, Hong Kong

(February 1, 2008)

We show that any critical transition region between two adjacent Hall plateaus in either integer or fractional quantum Hall effect is characterized by a universal “semi-circle” relationship between the longitudinal and transverse conductivities σ_{xx} , σ_{xy} , provided the sample is homogeneous and isotropic on a large scale. This is demonstrated both for the phase-coherent quantum transport as well as for the incoherent transport.

PACS number: 73.40.Hm

Recently there has been much interest in the study of transition regions between successive quantum Hall plateaus, in both the integer and the fractional regime. Most of these studies have been focused on how the transition region width ΔB scales with temperature at low temperatures. Experimentally, it has been noted that the critical dependence $\Delta B \propto T^\kappa$ can be extracted either directly from the half-width of the peak in $\sigma_{xx}(B)$, or from the sharpness of the step in Hall conductivity $\sigma_{xy}(B)$. Thus it has been realized, on intuitive ground at least, that the peak-like $\sigma_{xx}(B)$ function and the step-like $\sigma_{xy}(B)$ function in the transition region between two successive plateaus are related. Based on a semiclassical model that the transition region, in the critical regime, is represented by a random mixture of two liquids with different quantized local Hall conductivities σ_1 and σ_2 , Dykhne and Ruzin [1] recently developed a phenomenological theory to show that σ_{xx} and σ_{xy} are related by a simple “semi-circle” relation:

$$\sigma_{xx}^2 + \left(\sigma_{xy} - \frac{\sigma_1 + \sigma_2}{2} \right)^2 = \left(\frac{\sigma_1 - \sigma_2}{2} \right)^2. \quad (1)$$

Physically, a finite *effective* longitudinal conductivity σ_{xx} at $T \rightarrow 0$ is related to the existence of singularities in the current distribution (“hot spots”) close to the percolation threshold. To derive relation (1), authors in Ref. [1] assumed a very weak scattering in the bulk and described it by a *local* diagonal conductivity $\sigma_{xx}^{\text{loc}} \ll \sigma_2 - \sigma_1$, defined on a scale shorter than the correlation length of the two-phase mixture. This assumption is essentially classical and does not apply to the more experimentally relevant

low-temperature regime where quantum coherence exists on a much larger scale [1].

In this letter, we demonstrate that the same simple relationship between σ_{xx} and σ_{xy} holds in the quantum coherent transport regime as well. We achieve this by first proving a theorem that the semi-circle relation between σ_{xx} and σ_{xy} is equivalent to the perpendicularity between the average current densities in the two phases. We then show by simulations on the quantum percolation model that the currents are indeed perpendicular to a high accuracy, thus finishing the demonstration.

As in Ref. [1], let us consider two competing quantum Hall liquids in the presence of a long-range random potential, with the magnetic field determining volume fractions (close to 1/2), as depicted in Fig. 1(a). The “white” regions represent the phase with (quantized) Hall conductivity σ_1 , such that if this phase percolates freely throughout the sample, the system will be on the Hall plateau with $\sigma_{xy} = \sigma_1$. Similarly, the “black” regions have local Hall conductivity σ_2 . Near the percolation threshold, transport is controlled by quantum tunneling through the various saddle points (vertices), one of which is shown in detail in Fig. 1(b). In the presence of an external electric field, the non-equilibrium Hall currents flow inside the white and black regions. (Relation of this “bulk” transport description to more conventional Landauer-Büttiker formalism will be discussed later on). We assume that temperature is so low that scattering inside the bulk regions is absent. In this case, due to the continuity conditions, the currents are not allowed to cross the edges and must flow along them focusing at the saddle point [1]. Let us define the currents entering and leaving the white quadrants I and I' and those in the black quadrants J and J' as shown in Fig. 1(b). For any distribution of the current densities across the quadrants, one can relate the net currents to the four electric potentials on edges V_1 through V_4 , as given by

$$\begin{aligned} I &= \sigma_1(V_2 - V_1), \quad I' = \sigma_1(V_3 - V_4); \\ J &= \sigma_2(V_3 - V_2), \quad J' = \sigma_2(V_4 - V_1). \end{aligned} \quad (2)$$

As one can easily check, current conservation is met only if $V_2 - V_1 = V_3 - V_4$. Thus we have in fact $I = I'$ and $J = J'$, i.e. there is a unique white-to-white current and a unique black-to-black one for each saddle point.

The pattern of the current distribution in the sample is determined by the “white” and “black” currents I_i

and J_i , for each vertex i in the sample (Fig. 1(a)). This in general can be quite complex. Let $\vec{i}$ and $\vec{j}$ be the average partial current densities for the white and black phases respectively. The total current density is written $\vec{j}_{\text{tot}} = \vec{i} + \vec{j}$. The central rigorous result in this paper is the following theorem:

Theorem: If and only if $\vec{i} \perp \vec{j}$, the effective conductivity tensor $\hat{\sigma}$ obeys the semi-circle relation in Eq. (1). This theorem reduces the semi-circle relationship between σ_{xx} and σ_{xy} to a purely geometrical property of current distributions in the two phases.

Proof: Let us introduce, in addition to the partial current densities $\vec{i}$ and $\vec{j}$, the average “white” and “black” electric fields $\vec{e}$ and $\vec{f}$, related to the former by

$$\vec{i} = \sigma_1 \hat{e} \cdot \vec{e}, \quad \vec{j} = \sigma_2 \hat{e} \cdot \vec{f}, \quad (3)$$

where $\hat{e}$ is the matrix of clockwise rotation by 90° . These relations are valid only if scattering is absent everywhere except at separated points (vertices). One can obtain them, e.g., by drawing a vertical line across the sample as shown in Fig. 1(a). Adding total voltage drop along it, and the total current across it, for white and black regions separately, one obtains $i_x = \sigma_1 e_y$ and $j_x = \sigma_2 f_y$. Repeating this procedure for a horizontal line yields the remaining relations in Eqs. (3). Now suppose $\vec{i}$ and $\vec{j}$ are perpendicular to each other. This can be written as

$$\vec{e} = \beta \hat{e} \cdot \vec{f}, \quad (4)$$

where β is some dimensionless parameter. The total current density and field are then given by $\vec{j}_{\text{tot}} = \vec{i} + \vec{j}$, $\vec{e}_{\text{tot}} = \vec{e} + \vec{f}$. Using Eqs. (4) and (3), we can relate $\vec{j}_{\text{tot}}$ to $\vec{e}_{\text{tot}}$ directly, $\vec{j}_{\text{tot}} = \hat{\sigma}_{\text{eff}} \cdot \vec{e}_{\text{tot}}$, where the effective conductivity tensor $\hat{\sigma}_{\text{eff}}$ is given by

$$\hat{\sigma}_{\text{eff}} = \frac{\sigma_1 + \sigma_2}{2} \hat{e} + \frac{\sigma_1 - \sigma_2}{2} \hat{e} \frac{\beta + \hat{e}}{\beta - \hat{e}}. \quad (5)$$

It is easy to show that Eq. (5) implies

$$\det \left[\hat{\sigma}_{\text{eff}} - \hat{e} \frac{\sigma_1 + \sigma_2}{2} \right] = \frac{(\sigma_1 - \sigma_2)^2}{4}, \quad (6)$$

independent of the value of β . This is equivalent to the semi-circle relation (1). It is also easy to see that if the condition $\vec{i} \perp \vec{j}$ is not satisfied, we would have $\hat{\beta} = \beta \hat{\Gamma}$, where $\hat{\Gamma}$ is some rotation matrix. In this case, the determinant in Eq. (6) would depend on β . Thus $\vec{i} \perp \vec{j}$ is also a necessary condition for the semi-circle relation (1). Q.E.D.

We now verify analytically that $\vec{i} \perp \vec{j}$ holds for an incoherent (classical) system at high T . This regime is characterized by a relatively short phase-coherence length, $l_\phi \ll \xi_p$, where ξ_p is the percolation correlation length determining the typical distance between vertices. In this case, scattering at a vertex has a local nature, the black

and white currents at each vertex i are uniquely related by

$$I_i = \alpha_i J_i, \quad (7)$$

where parameter α_i depends on the scattering properties of this vertex [1]. For our purposes, it suffices to consider this quantity as a phenomenological parameter which fluctuates from vertex to vertex. As one can prove, Eq. (7) and the condition that the sum of currents entering each (black or white) region must be zero, uniquely define all the currents $\{I_i, J_i\}$, given the average current density $\vec{j}_{\text{tot}}$. Assuming that chirality is absent in the neighborhood of all the saddle points, parameters α_i must change sign when the magnetic field is reversed

$$\alpha_i(B) = -\alpha_i(-B). \quad (8)$$

We now show that this condition leads to $\vec{i} \perp \vec{j}$. Let $\{I_i, J_i\}$ be the current distribution for some magnetic field B and some $\vec{j}_{\text{tot}}$, with $\vec{i}$ and $\vec{j}$ being the partial current densities. At the opposite direction of the magnetic field, $B' = -B$, there exists a solution $\{I'_i, J'_i\}$ such that $I'_i = I_i$ and $J'_i = -J_i$. The partial current densities are $\vec{i}' = \vec{i}$ and $\vec{j}' = -\vec{j}$. On the other hand, since the system is isotropic and non-chiral at large length scales, the pair of vectors $\vec{i}, \vec{j}'$ can be obtained from the pair $\vec{i}, \vec{j}$ by rotation and reflection. These conditions are compatible only if $\vec{i} \perp \vec{j}$. The proof given can be easily generalized to the more complex case where each separate vertex has a random chirality, but the system as a whole is non-chiral.

Before going on to show $\vec{i} \perp \vec{j}$ in the quantum transport regime, let us discuss a rather general issue of the relation between bulk and edge-state transport pictures when applied to our problem. The bulk current description employed above [2]- [4] differs from the conventional edge-state formalism which assumes that current flows directly on the edge. This discrepancy comes from the use of single-electron picture in the latter theory: non-equilibrium current in the bulk is impossible since both concentration of electrons and built-in electric field (in our case, that from random potential) are fixed. Adding electrons to the edge by shifting chemical potentials results in this picture only in an additional edge current. But since real electrons do have charge, accumulation of electrons at edge-states (and of localized quasiparticles in the bulk) generates an electric field $\vec{E}(\vec{r})$ inside the bulk regions in Fig. 1(a), causing additional Hall current $\vec{j}(\vec{r})$ to flow in the bulk [5]. Using Lorentz transformation, one can show that a homogeneous interacting electron system subject to a homogeneous electric field acquires a drift velocity equal to the classical value for a single electron. In the spirit of linear response, it is natural to assume that the same is true locally for a weakly inhomogeneous case: $\vec{j}(\vec{r}) = \sigma_{xy}(\vec{r}) \hat{e} \cdot \vec{E}(\vec{r})$, where $\sigma_{xy}(\vec{r}) = en(\vec{r})c/B$. Our next assumption is that quasi-particles in the bulk

are localized by short-range impurities, so that the effective concentration n which contribute to the Hall current is given by that of the incompressible liquid. As a result, integrating electric field and the current density along any curve connecting the opposite sides of an angle in Fig. 1(b) gives relations of Eqs. (2).

Although the above arguments suggest that the local σ_{xy} picture employed here is more appropriate for describing current distributions in the case of long-range electron interaction than the conventional edge-state formalism, it turns out that, in fact, both descriptions will yield identical results, in the absence of scattering in the bulk. This is because, in the edge-state formalism, the current density can be written as

$$\vec{j}_{\text{edge}} = \varphi(\vec{r}) \hat{e} \cdot \nabla \sigma_{xy}, \quad (9)$$

where $e\varphi(\vec{r})$ is the chemical potential function. In the local σ_{xy} picture, the corresponding current density is given by

$$\vec{j}_{\text{loc}} = -\sigma_{xy} \hat{e} \cdot \nabla \varphi(\vec{r}), \quad (10)$$

where $\varphi(\vec{r})$ is now the electric potential function. For step-like distribution of σ_{xy} [from white region to black region in Fig. 1(a)], Eq. (9) yields a delta-function on the edge with an integrated current $(\sigma_2 - \sigma_1)\varphi_{\text{edge}}$. As is easy to check, the difference between the two current densities (9), (10) is a curl which yields zero upon integration along any line throughout the whole sample, when the two φ functions in the two formalisms are identified to one another. Thus we are free to choose either formalism according to our convenience in our problem.

We now proceed to show that $\vec{i} \perp \vec{j}$ for coherent quantum percolation systems. At very low temperatures, different clusters of the same liquid couple to each other through quantum tunneling to form a coherent wavefunction whose size is larger than the classical cluster size ξ_p and diverges when approaching the percolation threshold. Quantum coherence implies that currents at a given saddle point can depend on those at other ones, so that the local relation Eq. (7) is no longer valid. Moreover, it is no longer sufficient to use currents $\{I_i, J_i\}$ to describe a given state of the system, since complex quantum amplitudes must be used instead. Thus we need to resort to a particular quantum percolation model. Our choice is the fully quantum coherent network model first suggested by Chalker and Coddington [6], which has proven quite successful in the study of localization length exponent in quantum Hall systems [6], [7]. In this model, a complex amplitude A_k is assigned to each bond of an oriented regular square network [see Fig. 1(c)]. At the vertex marked Ω in Fig. 1(c), the four quantum amplitudes are related by a scattering matrix given by

$$\begin{pmatrix} A_4 \\ A_3 \end{pmatrix} = \begin{pmatrix} e^{i\phi_4} & 0 \\ 0 & e^{i\phi_3} \end{pmatrix} \begin{pmatrix} \cosh \gamma \sinh \gamma \\ \sinh \gamma \cosh \gamma \end{pmatrix} \begin{pmatrix} e^{i\phi_1} & 0 \\ 0 & e^{i\phi_2} \end{pmatrix} \begin{pmatrix} A_1 \\ A_2 \end{pmatrix} \quad (11)$$

where ϕ_k are random phase factors which model the effect of random path lengths in an actual random geometry such as in Fig. 1(a), γ is the tunneling parameter which, for the sake of simplicity, is set to be the same for all the vertices. For vertices oriented like the one marked Γ in Fig. 1(c), the bonds denoted $A_1, \dots, A_4$ should be rotated by 90° with respect to those with orientation Ω . The limits $\gamma \rightarrow 0$ and $\gamma \rightarrow \infty$ correspond to 100% percolation of the white or the black regions, respectively. The value $\gamma_c = \ln(1 + \sqrt{2})$ defines the quantum percolation threshold [6]. The system of Eqs. (11) written for all the saddle points determines the distribution of the quantum amplitudes, i.e. the wavefunction.

In order to complete the construction of our formalism, we need to adapt this model to the study of transport properties. Original quantum percolation model [6] used the edge-transport theory as a starting point, its classical analog being the “network model” suggested in Ref. [8]. The squares of amplitudes $|A_k|^2$ were identified with the edge currents entering and leaving the vertices as shown in Fig. 1(c). Unitarity of the scattering matrix in Eq. (11) takes care of current conservation: $|A_1|^2 - |A_2|^2 = |A_4|^2 - |A_3|^2$. Obviously, the standard edge-transport theory cannot be applied consistently to the quantum case since the chemical potential it deals with cannot be introduced for separate bonds belonging to the same wavefunction. Hence we identify $|A_k|^2$ with the fictitious edge current $(\sigma_2 - \sigma_1)V_k$ discussed above in a formal sense, with V_k being the *electric* potential of the bond. Then, as one can see from Eqs. (2), the physical currents in the white and black quadrants [Fig. 1(b)] are related to the differences in the edge currents on the adjoining edges, given by

$$I = \sigma_1 (|A_2|^2 - |A_1|^2), \quad J = \sigma_2 (|A_3|^2 - |A_4|^2), \quad (12)$$

apart from a constant factor $\sigma_1 - \sigma_2$. Since we study small non-equilibrium currents everywhere in this work, Eqs. (12) can be linearized

$$\begin{aligned} I &= \sigma_1 [\text{Re}(A_2 a_2^*) - \text{Re}(A_1 a_1^*)], \\ J &= \sigma_2 [\text{Re}(A_3 a_3^*) - \text{Re}(A_4 a_4^*)], \end{aligned} \quad (13)$$

where a_k are small perturbations of the quantum amplitudes due to an applied external voltage V , while A_k are the equilibrium values in the absence of V . Quantities $\text{Re}(A_k a_k^*)$ are the non-equilibrium electric potentials.

The remaining task is to solve the system of linear Equations (11) numerically with proper boundary conditions and find the average black and white current densities using Eqs. (13). We chose standard periodic boundary conditions on the left and right boundaries of a rectangular sample in Fig. 1(c). Since Eqs. (11) are linear, they can be solved separately for A_k and a_k , but with different boundary conditions. We let all equilibrium amplitudes A_k on the incoming bonds on the upper and lower boundaries to be unity. For the same sample,

we let incoming amplitudes a_k on the lower boundary to be $+V/2$ and those on the upper boundary to be $-V/2$. This simulates the situation where a small two-terminal voltage V is applied across the Corbino disc-shaped sample attached to two metallic contacts. Indeed, as shown by one of us when studying the role of contacts in the non-uniform Hall conductors [3], a potential step near the contact is possible if and only if local σ_{xy} increases to the right when looking into the sample from the contact. This is the case for the bonds arrowed as outgoing ones in Fig. 1(c) ($\sigma_1 < \sigma_2$ is implied). The incoming bonds, on the contrary, must have potentials of the contacts. This is completely analogous to the “ballistic” boundary conditions in the conventional edge-state transport theory where the incoming (leaving a contact) one-electron states are assumed to have the chemical potential of the contact.

For each realization of $\{\phi_k\}$, we can extract the currents densities $\vec{i}$ and $\vec{j}$ by adding up the corresponding currents at the vertices, according to Fig. 1(c). Namely, i_x and j_y are found by averaging I and J currents, respectively, over all the vertices with orientation Ω (Fig. 1(c)). Similarly, i_y and j_x are extracted from the vertices Γ . The total number of rows of vertices in either direction is even. The system size $L_x \times L_y$ ranges from 20×20 to 50×82 , and we averaged over 2,000 realizations to find the ensemble averaged $\langle \vec{i} \rangle$ and $\langle \vec{j} \rangle$ currents. We note the importance of ensemble averaging: in the fully quantum mechanical case studied here, there is no simple self-averaging even in a large sample. Self-averaging occurs, in a physical system, only on the length scale of an inelastic phase-breaking length, which at low temperatures can be very large, the effect of which is replaced in our simulation by a finite system size. Our result for $\gamma = \gamma_c = \ln(1 + \sqrt{2})$ can be summarized as

$$\cos \alpha = \frac{\langle \vec{i} \rangle \cdot \langle \vec{j} \rangle}{|\langle \vec{i} \rangle| |\langle \vec{j} \rangle|} = 0.01 \pm 0.02. \quad (14)$$

This shows that the white and black currents are indeed perpendicular to each other to a high degree of accuracy. We have checked that this null value for $\cos \alpha$ persists when the size of the sample is changed. Neither did we find any dependence of $\cos \alpha$ on the sample shape for any $L_y/L_x > 1.3$. Finally, we made sure that $\cos \alpha$, in such samples, is insensitive to changes of γ around the critical value γ_c . The interval γ in which the ratio of $|\vec{i}|/|\vec{j}|$ changes by 2 orders of magnitude was studied. (A weak shape dependence at $\gamma = \gamma_c$ was revealed in short samples: at $L_y/L_x = 0.3$, $\cos \alpha = 0.14 \pm 0.02$. We interpret this as an artifact of closely placed metallic contacts; since when the localization length becomes smaller than the sample size as we take $\gamma \neq \gamma_c$, $\cos \alpha$ restores its null value.) In combination with the Theorem discussed before, we have thus successfully demonstrated that the universal semi-circle relation between σ_{xx} and σ_{xy} holds

in the regime of quantum transport.

We thank F. Zeng for assistance in simulations, and S. Kivelson for many stimulating discussions. This work was supported in part by the ONR Grant N00014-90-J-1829, and the DOE Grant DE-FG03-88ER45378.

-
- [1] A.M. Dykhne and I.M. Ruzin, Phys. Rev. B, in print.
 - [2] R. Woltjer, R. Eppenga, and M.F.H. Schuurmans, in *High Magnetic Fields in Semiconductor Physics*, edited by G. Landwehr (Springer, Berlin, 1987), p. 104.
 - [3] I.M. Ruzin, Phys. Rev. B, Phys. Rev. B **47**, 15727 (1993).
 - [4] N.R. Cooper and J.T. Chalker, Phys. Rev. B **48**, 4530 (1993).
 - [5] D.J. Thouless, Phys. Rev. Lett. **71**, 1879 (1993).
 - [6] J.T. Chalker and P.D. Coddington, J. Phys. C **21**, 2665 (1988).
 - [7] D.-H. Lee, Z. Wang, and S. Kivelson, Phys. Rev. Lett. **70**, 4130 (1993).
 - [8] J. Kucera and P. Streda, J. Phys. C **21**, 4357 (1988).

FIG. 1. (a) Schematic illustration of the random quantum percolation model for describing the transition between successive quantum Hall plateaus. (b) A single saddle point magnified. (c) The Chalker-Coddington network model ($\sigma_1 < \sigma_2$).